

Package Leaflet:
Information for the patient

circadin® 2 mg
prolonged-release tablets
Melatonin

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See Section 4.

What is in this leaflet:

1. What Circadin is and what it is used for
2. What you need to know before you take Circadin
3. How to take Circadin
4. Possible side effects
5. How to store Circadin
6. Contents of the pack and other information

1. What Circadin is and what it is used for

The active substance of Circadin, melatonin, belongs to a natural group of hormones produced by the body. Circadin is used on its own for the short-term treatment of primary insomnia (persistent difficulty in getting to sleep or staying asleep, or poor quality of sleep) in patients aged 55 years and older. 'Primary' means that the insomnia does not have any identified cause, including any medical, mental or environmental cause.

2. What you need to know before you take Circadin

Do not take Circadin

- if you are allergic to melatonin or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions

- Talk to your doctor or pharmacist before taking Circadin.
- If you suffer from liver or kidney problems. No studies on the use of Circadin in people with liver or kidney diseases have been performed, you should speak to your doctor before taking Circadin as its use is not recommended.
 - If you have been told by your doctor that you have an intolerance to some sugars.
 - If you have been told you suffer from an autoimmune disease (where the body is 'attacked' by its own immune system). No studies on the use of Circadin in people with auto-immune diseases have been performed; therefore, you should speak to your doctor before taking Circadin as its use is not recommended.
 - Circadin can make you feel drowsy, you should be careful if the drowsiness affects you as it may impair your ability on tasks such as driving.
 - Smoking may make Circadin less effective, because the components of tobacco smoke can increase the breakdown of melatonin by the liver.

Children and adolescents

Do not give this medicine to children between the ages of 0 to 18 years as it has not been tested and its effects are unknown.

Other medicines and Circadin

- Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. These medicines include:
- Fluvoxamine (used for the treatment of depression and obsessive compulsive disorder), psoralens (used in the treatment of

skin disorders e.g. psoriasis), cimetidine (used in the treatment of stomach problems such as ulcers), quinolones and rifampicin (used in the treatment of bacterial infections), oestrogens (used in contraceptives or hormone replacement therapy) and carbamazepine (used in the treatment of epilepsy).

- Adrenergic agonists/antagonists (such as certain types of medicines used to control blood pressure by constricting blood vessels, nasal decongestants, blood pressure lowering medicines), opiate agonists/antagonists (such as medicinal products used in the treatment of drug addiction), prostaglandin inhibitors (such as nonsteroidal anti-inflammatory medicines), antidepressant medication, tryptophan and alcohol.
- Benzodiazepines and non-benzodiazepine hypnotics (medicines used to induce sleep such as zaleplon, zolpidem and zopiclone)
- Thioridazine (for the treatment of schizophrenia) and imipramine (for the treatment of depression).

Circadin with food, drink and alcohol

Take Circadin after you have eaten. Do not drink alcohol before, during or after taking Circadin, because it reduces the effectiveness of Circadin.

Pregnancy and breast-feeding

Do not take Circadin if you are pregnant, think you may be pregnant, trying to become pregnant or breast-feeding. Ask your doctor or pharmacist for advice before taking this medicine.

Driving and using machines

Circadin may cause drowsiness. If you are affected, you should not drive or operate machinery. If you suffer from continued drowsiness, then you should consult your doctor.

Circadin contains lactose monohydrate

Circadin contains lactose monohydrate. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

3. How to take Circadin

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose is one Circadin tablet (2 mg) taken daily by mouth, after food, 1-2 hours before bedtime. This dosage may be continued for up to thirteen weeks. You must talk to a doctor if you do not feel better or if you feel worse after 21 days.

You should swallow the tablet whole. Circadin tablets should not be crushed or cut in half.

If you take more Circadin than you should

If you have accidentally taken too much of your medicine, contact your doctor or pharmacist as soon as possible.

Taking more than the recommended daily dose may make you feel drowsy.

If you forget to take Circadin

If you forget to take your tablet, take another as soon as you remember, before going to sleep, or wait until it is time to take your next dose, then go on as before.

Do not take a double dose to make up for a forgotten dose.

If you stop taking Circadin

There are no known harmful effects if treatment is interrupted or ended early. The use of Circadin is not known to cause any withdrawal effects after treatment completion.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can

cause side effects, although not everybody gets them.

If you experience any of the following serious side effects, stop taking the medicine and contact your doctor **immediately:**

Uncommon: (may affect up to 1 in 100 people)
– Chest pain

Rare: (may affect up to 1 in 1000 people)

- Loss of consciousness or fainting
- Severe chest pain due to angina
- Feeling your heartbeat
- Depression
- Visual impairment
- Blurred vision
- Disorientation
- Vertigo (a feeling of dizziness or "spinning")
- Presence of red blood cells in the urine
- Reduced number of white blood cells in the blood
- Reduced blood platelets, which increases risk of bleeding or bruising
- psoriasis

If you experience any of the following non-serious side effects contact your doctor and/or seek medical advice:

Uncommon: (may affect up to 1 in 100 people)

Irritability, nervousness, restlessness, insomnia, abnormal dreams, nightmares, anxiety, migraine, headache, lethargy (tiredness, lack of energy), restlessness associated with increased activity, dizziness, tiredness, high blood pressure, upper abdominal pain, indigestion, mouth ulceration, dry mouth, nausea, changes in the composition of your blood which could cause yellowing of the skin or eyes, inflammation of the skin, night sweats, itching, rash, dry skin, pain in extremities, menopausal symptoms, feeling of weakness, excretion of glucose in the urine, excess proteins in the urine, abnormal liver function and weight increase.

Rare: (may affect up to 1 in 1000 people)

Shingles, high level of fatty molecules in the blood, low serum calcium levels in the blood, low sodium levels in the blood, altered mood, aggression, agitation, crying, stress symptoms, early morning awakening, increased sex drive, depressed mood, memory impairment, disturbance in attention, dreamy state, restless legs syndrome, poor quality sleep, 'pins and needles' feeling, watery eyes, dizziness when standing or sitting, hot flushes, acid reflux, stomach disorder, blistering in the mouth, tongue ulceration, stomach upset, vomiting, abnormal bowel sounds, wind, excess saliva production, bad breath, abdominal discomfort, gastric disorder, inflammation of the stomach lining, eczema, skin rash, hand dermatitis, itchy rash, nail disorder, arthritis, muscle spasms, neck pain, night cramps, prolonged erection that might be painful, inflammation of the prostate gland, tiredness, pain, thirst, passing large volumes of urine, urinating during the night, increased liver enzymes, abnormal blood electrolytes and abnormal laboratory tests.

Frequency not known: (cannot be established from the available data)
Hypersensitivity reaction, swelling of mouth or tongue, swelling of the skin and abnormal milk secretion.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via:

United Kingdom

Yellow Card Scheme
Website: www.mhra.gov.uk/yellowcard

Ireland

Pharmacovigilance Section
Irish Medicines Board
Kevin O'Malley House
Earlsfort Centre, Earlsfort Terrace
IRL - Dublin 2

Tel: +353 1 6764971
Fax: +353 1 6762517
Website: www.imb.ie
e-mail: imbpharmacovigilance@imb.ie

Malta

ADR Reporting
The Medicines Authority
Post-Licensing Directorate
203 Level 3, Rue D'Argens
GZR-1368 Gzira
Website:
www.medicinesauthority.gov.mt
e-mail:
postlicensing.medicinesauthority@gov.mt

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Circadin

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton (EXP). The expiry date refers to the last day of that month.

Do not store above 25°C. Store in the original package in order to protect from light.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Circadin contains

- The active substance is melatonin. Each prolonged-release tablet contains 2 mg melatonin.
- The other ingredients are ammonio methacrylate copolymer type B, calcium hydrogen phosphate dihydrate, lactose monohydrate, silica (colloidal anhydrous), talc and magnesium stearate.

België/Belgique/Belgien, Luxembourg/Luxemburg

Takeda Belgium
Chaussée de Gand 615 Gentessteenweg
B-1080 Bruxelles / Brussel / Brüssel
Tél./Tel: +32 2 464 06 11
Takeda-Belgium@takeda.com

България, Česká republika, Hrvatska, Κύπρος, Magyarország, Malta, România, Slovenija, Slovenská republika

RAD Neurim Pharmaceuticals EEC Limited
One Forbury Square
The Forbury
Reading
Berkshire RG1 3EB
United Kingdom
Tel: +44 7563543352
neurim@neurim.com

Danmark, Island

Takeda Pharma A/S
Dybendal Alle 10,
DK-2630 Taastrup
Danmark
Tlf./Simi: +45 46 77 11 11
info@Takeda.dk

Deutschland

Medice Arzneimittel Pütter GmbH & Co. KG
Kuhloweg 37
D58638 Iserlohn
Tel: +49 (0)2371 9370
info@medice.de

Eesti

Takeda Pharma AS
Jaama 55b
EE-63308 Põlva
Tel: +372 6112 569

Ελλάδα

TAKEDA ΕΛΛΑΣ Α.Ε.
Λεωφ. Κηφισίας 196
152 31 Χαλάνδρι, Αθήνα
Τηλ: +30 210 6729570
gr.info@takeda.com

What Circadin looks like and contents of the pack

Circadin 2 mg prolonged-release tablets are available as white to off-white round biconvex shaped tablets. Each carton of tablets contains one blister strip of 7, 20 or 21 tablets, or alternatively in a carton containing two blister strips of 15 tablets each (30 tablet pack). Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder:
RAD Neurim Pharmaceuticals EEC Limited
One Forbury Square
The Forbury
Reading
Berkshire RG1 3EB
United Kingdom
e-mail: neurim@neurim.com

Manufacturer:

Sites responsible for Batch Release in the EEA:
Temmler Pharma GmbH & Co. KG
Temmlerstrasse 2
35039 Marburg
Germany

Iberfar - Indústria Farmacéutica S.A.
Rua Consiglieri Pedroso, n.o 121123 -
Queluz de Baixo
Barcarena, 2745557
Portugal

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder

This leaflet was last revised in 02/2014.

Other sources of information

Detailed information on this medicine is available on the European Medicines Agency web site: <http://www.ema.europa.eu>

España

JUSTE S.A.Q.F.
Avda. San Pablo, 27
28823 Coslada
Madrid
Tel.: + 34 91 724 82 00
medicocientifica@juste.net

France

BIOCODEX
7 Avenue Gallieni
94150 Gentilly
Tel: +33 (0)1 41 24 30 00
webar@biocodex.fr

Ireland, United Kingdom

Flynn Pharma Limited
Hertlands House
Primett Road
Stevenage
Hertfordshire SG1 3EE
United Kingdom
Tel: +44 (0) 1438 727822
medinfo@flynnpharma.com

Italia

Fidia Farmaceutici S.p.A
Via Ponte della Fabbrica 3/A
35031 Abano Terme (PD)
Tel: +39 049 8232355
Fax: +39 049 810653
info@fidiafarma.it

Latvija

Takeda Latvia SIA
Duntes iela 6
Rīga, LV 1013
Tel: +371 784 0082

Lietuva

Takeda, UAB
Gynėjų 16
LT-01109 Vilnius
Tel: +370 521 09 070
lt-info@takeda.com

Nederland

Takeda Nederland bv
Jupiterstraat 250
2132 HK Hoofddorp
Tel: +31 23 566 8777
nl.medical.info@takeda.com

Norge

Takeda Nycomed AS
Drammensveien 852
N-1372 Asker
Tel: +47 6676 3030
infor norge@takeda.com

Österreich

SANOVA PHARMA GesmbH
Haidestraße 4
1110 Wien
Tel.: +43 (01) 80104-0
Fax: +43 (01) 8042904
sanova.pharma@sanova.at

Polska

MEDICE Arzneimittel Pütter GmbH & Co. KG
ul. Chorzowski Pancerniej 27
02-951 Warszawa
Tel.: + 48-(0)22 642 2673
office@medice.pl

Portugal

Italfarmaco, Produtos Farmacéuticos, Lda.
Rua Consiglieri Pedroso, 123
Queluz de Baixo
2730-056 BARCARENA, Portugal
Tel. +351 214 342 530
geral@itf-farma.pt

Suomi/Finland

Oy Leiras Takeda Pharmaceuticals Ab
PL/PB 1406
FIN-00101 Helsinki
Puh./Tel: +358 20 746 5000
info@leiras.fi

Sverige

Takeda Pharma AB
Box 3131, 169 03 Solna
Tel: +46 8 731 28 00
infosweden@takeda.com

Important

Please ensure that the **TEXTS** and **LAYOUTS** are to your satisfaction as *Rami Raviv Graphic Design Ltd.* can not accept liability for any errors overlooked once this proof has been signed and returned.

All **BAR CODES** and **BRILLE TEXT** must be verified by the client/manufacturer/printer. *Rami Raviv Graphic Design Ltd.* is unable to do this and can not accept liability for any errors.

COLORS may not reproduce accurately on this proof. You should refer to Pantone Reference Guides for accurate color match.

Date	Reason for Draft
20/08/2013	New Annex, Sept 2013
15/01/2014	Iberfar, Partners Address, 02/14

Product Title	CIRCADIN	Date	15/01/2014
Strength	2 mg	Client	Neurim Pharma
Item	Patient Info Leaflet	Language	EN
Format	170 X 400 mm	Country	EN / IR / MA
Typefaces	Myriad Pro	Type Size	9 pt / 10 pt
No of Colours	1	Job Name	PIL_ENIRMA_01
			Black

Rami Raviv Graphic Design Ltd. • +972-3-6969838 • ravivdmg@netvision.net.il